

DRAFT REPORT ON
COASTAL ZONE MANAGEMENT PLAN (CZMP) FOR
THRISSUR DISTRICT – KERALA
(Prepared as per CRZ Notification 2019)



Prepared for
Department of Environment
Government of Kerala



NATIONAL CENTRE FOR EARTH SCIENCE STUDIES
Ministry of Earth Sciences, Government of India
Thiruvananthapuram - 695 011, Kerala

March 2023

DRAFT REPORT ON
COASTAL ZONE MANAGEMENT PLAN (CZMP) FOR
THRISSUR DISTRICT – KERALA
(Prepared as per CRZ Notification 2019)

Prepared for
Department of Environment
Government of Kerala



NATIONAL CENTRE FOR EARTH SCIENCE STUDIES
Ministry of Earth Sciences, Government of India
Thiruvananthapuram - 695 011, Kerala

March 2023

CONTENTS

	Page No:
1: INTRODUCTION	1
1.1: CZMP Planning Process	4
1.2: Development of a coastal database and information system	5
1.3: Generation of CZMP maps	6
2: THE STATE OF KERALA	7
2.1: Thrissur District	9
2.1(a): Administration	9
2.1(b): Physiography	10
2.1(c): Rivers and drainage characteristics	12
2.1(d): Coastal Wetlands	14
2.2: Demography and socio-economic activities	15
2.3: Coastal Geomorphology and Ecosystem	16
2.4: Marine Fishery Resources	17
2.5: Biodiversity of Thrissur district	19
2.6: Pollution and Waste Management Issues	20
3: PURPOSE AND SCOPE OF CZMPs	21
4: COASTAL ZONE MANAGEMENT PLANS	21
4.1: Demarcation of High Tide Line (HTL) and Low Tide Line (LTL)	22
4.1 (a): Landward (monsoonal) berm crest for beaches	24
4.1 (b): Seawall/revetments/embankments	24
4.1 (c): Permanent Vegetation Line	24
4.1 (d): Coastal sand dune / paleo-aeolian dune	25
4.1 (e): Mangroves	25
4.1 (f): Rocks, Headlands, Cliffs	29
4.1 (g): Influence of Tidal action	29
4.2: Demarcation of Ecologically Sensitive Areas	30
5: ECOLOGICALLY SENSITIVE AREAS/COASTAL LANDUSE	31
5.1: Mangroves	31
5.2: Coral Reefs	32
5.3: Reserve Forests	32
5.4: Sand Dunes	32
5.5: Salt Marsh	32
5.6: Nesting Ground of Birds	32
5.7: Archaeologically important and Heritage Sites	32

5.8:	Seagrass	34
5.9:	Mudflats	34
5.10:	Turtle Nesting Grounds	34
5.11:	Inter-Tidal Zone	34
5.12:	Salt pan / Aquaculture ponds	35
6:	METHODOLOGY FOR PREPARATION OF CZMP	35
6.1:	Field Mapping and Map Preparation	36
7:	CRZ CLASSIFICATION	37
7.1:	CRZ - 1	37
	7.1.1: CRZ - 1 A	37
	7.1.2: CRZ - 1 B	38
7.2:	CRZ - II	38
7.3:	CRZ - III	38
	7.3.1: CRZ - III A	38
	7.3.2: CRZ - III B	38
7.4:	CRZ - IV	39
	7.4.1: CRZ - IV A	39
	7.4.2: CRZ - IV B	39
7.5:	Regulation limits/lines	39
7.6:	CVCA and IIIMP	40
8:	HAZARD LINE	41
9:	CRZ CATEGORIES OF THRISSUR DISTRICT	41
9.1:	CRZ Categories and ESAs (Panchayat/Village-wise) in Thrissur District	42
10:	CONCLUSION	42
ANNEXURE		
Annexure-1:	MAPS:	43
	MapNo.1: An overview of the distribution of CRZ Area among local bodies- Thrissur	44
	MapNo.2: Map grids with their numbers covering Thrissur District (12 map frames bearing Map Numbers KL-37, KL-38, KL-39, KL-40, KL-41, KL-42, KL-43, KL-44, KL-45, KL-46, KL-47, KL-48)	45
	MapNo.3: Location map of Thrissur District, Kerala	46
	MapNo.4: Overall distribution of Mangroves in Thrissur District	47
	MapNo.5: Archeological and Heritage sites in Thrissur District	48

MapNo.6: Inter Tidal Zone in Thrissur District	49
MapNo.7: Locations of turtle nesting grounds along the Thrissur District coast	50
MapNo.8: The Coastal villages categorized as CRZ- III A and CRZ III B in Thrissur District	51
Annexure-2: TABLES	52
Tables - 1: Comparison of CZMP 2011 and draft CZMP based on CRZ Notification 2019	53
Tables - 2: Village-wise population statistics for identifying the CRZ-III A category - Thrissur district	54
Tables - 3: CRZ Details in Local bodies of Thrissur	55
Tables - 4: CRZ Details in Villages of Thrissur District	57
Tables - 5: List of backwater and main coast islands in Thrissur District	60
Annexure-3: FIELD PHOTOGRAPHS	62
Plate - 1: HTL at Chettuva Azhimukham beach. (Location: 76° 2' 31.15" E 10° 30' 13.47" N, Vadanappally Panchayat, Village: Engandiyoor)	63
Plate - 2: HTL at Nakshathra beach (Location: 76° 3' 7.97" E 10° 29' 5.55" N, Vadanappally Panchayat, Village: Vadanappally)	63
Plate - 3: Mangrove area near William Fort Chettuva, (Location: 76° 2' 13.10" E 10° 31' 34.31" N, Engandiyoor Panchayat, Village: Engandiyoor)	64
Plate - 4: Mangrove patch near William Fort, Kotta Road (Location: 76° 2' 16.57" E 10° 31' 33.09" N, Engandiyoor Panchayat, Village: Engandiyoor)	64
Plate - 5: HTL at Munakkakadavu, Kadappuram (Location: 76° 2' 19.16" E 10° 30' 50.15" N, Orumanayur Panchayat, Village: Kadappuram)	65
Plate - 6: Mangrove area near KAMCO, Neythikudi bridge road (Location: 76° 14' 59.57" E 10° 14' 37.26" N, Mala Panchayat, Village: Vadama)	65
Plate - 7: Mangrove area at Kombathukadavu (Location: 76° 14' 16.80" E 10° 13' 53.27" N, Puthenchira Panchayat, Village: Puthenchira)	66
Plate - 8: Mangrove area near Poyya. (Location: 76° 14' 0.26" E 10° 12' 40.04" N, Poyya Panchayat, Village: Poyya)	66
Plate - 9: Turtle nesting site at Kadappuram Beach, Thottapu (Location: 76° 0' 50.21" E 10° 33' 36.55" N, Orumanayur Panchayat, Village: Kadappuram)	67

Plate - 10: Turtle nesting site at Edakkazhiyur beach (Location: 75° 59' 30.65" E 10° 36' 20.11" N, Chavakkad Municipality, Village: Manathala) 67

Annexure-4: OWNERSHIP DETAILS OF MANGROVES MORE THAN 1000SQ.M SPREAD AREA PROVIDED BY GOVT. OF KERALA

Annexure-5: SUMMARY OF TOURISM PLAN FOR THE CRZ AREA IN THRISSUR DISTRICT FORWARDED BY KCZMA.

Annexure-6: INTEGRATED FISHERIES DEVELOPMENT PLAN FOR CZMP, KERALA

COASTAL ZONE MANAGEMENT PLAN (CZMP) FOR THRISSUR DISTRICT - KERALA

1. INTRODUCTION

Coastal zone is the transitional zone, where land meets the sea and is influenced by both terrestrial and marine components. Intense interaction characterizes the coastal zone which varies from open sea to semi closed (creeks, lagoons) coastal waters and they exhibit a substantial diversity in environmental and demographic features. Here, land and ocean-dominated global processes converge and interact, characterized by multiple biogeochemical environmental gradients. The balance of these relationships provides a distinct domain of gradient-dependent ecosystems, climate, geomorphology, human habitation and most crucially regimes of highly dynamic physical, chemical and biological processes. Terrestrial processes are primarily governed by hydrological regimes and horizontal flows which provide mechanisms for energy gradients and transfer of materials (nutrients, contaminants, sediments), offering a range of conditions for material transformations and biological sustenance. Oceanic processes are similarly dominated by hydrological and physical factors that regulate the transit of materials and energy regimes, often in contrast with the land-dominated factors. The resultant equilibrium of terrestrial and oceanic processes yields regional and local heterogeneity in physical and ecological structure and supports the dynamics of ecosystem function and biogeochemical cycling in the coastal domain. Thus, mass and energy are constantly exchanging and as a result of these interactions created a unique ecosystem (Shailesh Nayak, 2017).

Coastal zones throughout the world have historically been among the most heavily exploited areas because of their rich resources. Coastal regions provide fish, shellfish, seaweeds and host ports for trading and commerce. In addition, several biota are sources of fertilizer, drugs, cosmetics and household products. Moreover, coastal wetlands also store and cycle nutrients, filter pollutants and help in the protection of the shoreline against erosion and storms. Thus, the richness and diversity of resources found in coastal regions have led to a corresponding concentration of human activities

and settlement along coasts and estuaries throughout the world. In coastal countries today an estimated half of the total populations live in coastal zones and migration from inland areas to the coast is increasing. Not surprisingly, there is also a sharp conflict between the need for immediate consumption or use of coastal resources and the need to ensure the long-term supply of those resources. In many countries this conflict has already reached a critical stage, with large parts of the coastal zone polluted from local or upland sources, fisheries severely degraded or destroyed, wetlands drained, coral reefs dynamited and beaches long since ruined for human enjoyment. If these coastal resources are to be maintained and restored, effective action is urgently needed. It is also obvious that the coastal zone will be expected to sustain the livelihoods of a very large proportion of the human population and will remain an important asset to people worldwide for the foreseeable future. The sustainability of the coastal environment is continuously impacted by pollution, eutrophication, industrialization, urban development, land reclamation, agricultural production, overfishing and exploitation. Moreover, the poor understanding of the dynamics of land-ocean interactions, coastal processes and the impact of poorly planned and managed human interventions makes the sustainability of human economic and social progress vulnerable to natural and human-induced hazards. Humans are increasingly influencing these regions, which resulting in measurable changes directly within the coastal domain and through feedback, indirectly within the terrestrial, oceanic and atmospheric compartments of the Earth system (Steffen et al., 2004). So, the major challenge that humans face today is how to manage the use of this area so that future generations can also enjoy its visual, cultural and societal resources. We need to ensure robust health of coastal ecosystems through sustainable management, so that they continue to provide various goods and services for future generations, as well.

According to a recent evaluation of the impacts of marine pollution from land-based sources, the degradation of the marine ecosystem is still occurring and, in many places has intensified (GESAMP, 2001). Hence, policies and legislations to reduce conflicts over uses in the coastal zone, protect coastal resources and support livelihood activities of local communities as well as to address the development requirements of the coast to meet economic and societal requirements are essential. Integrating environmental, economic and human activities to ensure pollution-free coastal waters and healthy ecosystems to sustain livelihood and coastal economy necessitates effective integration of science and public policy is very much needed. Due to various development schemes of private

and public, legal and illegal, large-scale modifications and damages to coastal morphology and ecosystems by way of reclamation of tidal flats, destruction of mangroves, leveling of sand dunes, mining of beach sand, construction activities for settlement, establishment of industries, dumping of waste and discharge of pollutants. Rapidly changing landuse due to the immense pressure for development in the coastal zone has adversely affected the coastal ecosystems, coastal morphology and livelihood resources of the coastal areas.

Coastal zone management depends on the information available on various aspects of coastal habitats, coastal processes, natural hazards and their impacts, water quality and living resources. The effective management techniques depend on such information and suitable response by concerned government agencies. Keeping these facts in view, Government of India on the recommendation of Ministry of Environment and forest (MoEF) passed a legislation called Coastal Regulation Zone (CRZ) in the year 1991. Under this legislative act, one of the most cost-effective long - term solutions to control various ecologically destructive activities in the endangered coastal zone, is to invoke spatial buffers around coastal ecosystems. The Coastal Regulation Zone (CRZ) Notifications (MoEF, 2019; 2011; 1991) provides buffer zones in the coastal area is being considered as the pragmatic tool to control, minimize and protect environmental damages to sensitive coastal stretches from unplanned human interference. Management of coastal ecosystems through CRZ requires identification and mapping of the regulation lines and the spatial extent of the ecosystems and morphologies in appropriate scales. Implementation and enforcement of the provisions of CRZ on the ground require extensive coastal mapping and continuous monitoring.

The National Centre for Earth Science Studies (NCESS), Thiruvananthapuram is an agency authorized by Government of India to prepare/update CZMP for the coastal stretches of our Country (OM F. No. J-17011/8/92-IA-III dated 08-08-2019). So, the Government of Kerala entrusted National Centre for Earth Science Studies (NCESS), Thiruvananthapuram for the preparation of Coastal Zone Management Plan (CZMP) for the State of Kerala following the guidelines in CRZ Notification 2019, vide G.O. (Rt) No. 80/2019/ENVT dated 28.08.2019. High Tide Line (HTL), Low Tide Line (LTL), Ecologically Sensitive Areas (ESAs) and Critically Vulnerable Coastal Areas (CVCAs) demarcated by the National Centre for Sustainable Coastal Management (NCSCM), Chennai, and the 'Hazard line' as demarcated by the Survey of India (SoI) have been made use for the preparation of CZMP.

1.1 CZMP Planning Process

The landmark Coastal Regulation Zone (CRZ) Notification, which was first issued on 19th February 1991, has been the most important legislative instrument in the country for coastal governance by considering the livelihood of fisherman and local people residing along the coast. The Ministry of Environment, Forests and Climate Change (MoEFCC), Government of India has issued a revised CRZ Notification on 6th January 2011 under Section 3(1) and Section 3(2)(v) of the Environment (Protection) Act, 1986 and Rule 5(3)(d) of Environment (Protection) Rules, 1986 in supersession of CRZ 1991 except as respect to things done or omitted to be done before such supersession. Subsequently, in June 2014, the MoEFCC constituted a committee under the chairpersonship of Dr. Shailesh Nayak, the erstwhile Secretary of the Ministry of Earth Sciences, to address the concerns raised by the state governments, eliminating ambiguities and simplifying certain provisions in CRZ Notification 2011. The committee held consultations with the state governments over the following six months and submitted its report to MoEFCC in January 2015. Ultimately, on 18th January 2019, in supersession of the CRZ Notification of 2011, the Government of India brought out the CRZ Notification 2019.

The CRZ is a critical regulation for conservation and livelihood protection on the coast. All developmental activities in the CRZ are regulated through the CRZ Notification. Accordingly, the CRZ has been declared as ‘the coastal stretches of the country and the water area up to its territorial water limit’. The Coastal Regulation Zone Notification (MoEF&CC, 2019; 2011; 1991) which provides buffer zones in the coastal area is being used as the best tool to control, minimize and protect environmental damages to sensitive coastal stretches from unplanned human interference. Thus, the CRZ Notification promote development in a sustainable manner based on scientific principles considering the dangers of natural hazards in the coastal areas and sea level rise due to global warming.

1.2 Development of a coastal database and information system

In the recent times, the availability of digital spatial data for the world coasts has vastly increased as a result of advancements in data capture and input techniques. The large increase in global data availability has had a significant impact on coastal science. The way in which coastal observations are stored and integrated largely determines the degree to which spatial processes can be understood. Therefore, well-organized and designed data systems are needed to underpin our understanding of the processes taking place over large parts of coasts. The expected accelerated rise in global mean sea levels may cause several physical changes to the world's coasts and hence can endanger coastal populations and infrastructure, as well as threaten many coastal ecosystems. The sensitivity of the coastal zone to sea-level rise, in conjunction with its importance in terms of social, economic and ecological value, highlights the need for consistent national- to global-scale assessments of potential impacts along the coasts. However, the scope of these studies has been limited by the available data in terms of resolution, coverage, parameter availability, and dated sources: this is a generic problem for broad-scale coastal analysis. In addition to these limitations, data quality and integration constitute further problems; even in those cases where data and tools are available to coastal scientists for the analysis and modeling of coastal processes, these usually exist in fragmented forms. This fact compromises the consistency, reliability and versatility of evaluations based on such sources. It has long been recognized that appropriate and reliable information within organized, planned and coherent coastal databases is an essential prerequisite for coastal zone management.

In order to address the preceding issues and provide a consistent source of data for the Indian coast, the data collated digitally onto a GIS platform for the preparation of the CZMP come handy. This database contains physical, ecological and vulnerability parameters and covers the Indian coasts uniformly, probably for the first time on a digital platform permitting retrieval, portability and sharing in a seamless manner. For this reason, the database has been specifically designed to address the data requirements of the project and the needs of researchers in the area of vulnerability assessment of coastal zones. It is also expected to be used for wider assessment of regional and global coastal issues.

1.3 Generation of CZMP maps

Management of coastal ecosystems through CRZ requires identification and mapping of the regulation lines and the spatial extent of the ecosystems and morphologies in appropriate scales. Implementation and enforcement of the provisions of CRZ on the ground require extensive coastal mapping and continuous monitoring. As per the CRZ Notification 2019 issued vide Notification No.G.S.R.37(E), dated the 18th January, 2019, all coastal States and Union territory administrations shall revise or update their respective coastal zone management plan (CZMP) framed under CRZ Notification, 2011 number S.O. 19(E), dated 6th January, 2011, as per provisions of this notification and submit to the Ministry of Environment, Forest and Climate Change for approval at the earliest and all the project activities attracting the provisions of this notification shall be required to be appraised as per the updated CZMP under this notification and until and unless the CZMPs is so revised or updated, provisions of this notification shall not apply and the CZMP as per provisions of CRZ Notification, 2011 shall continue to be followed for appraisal and CRZ clearance to such projects. The Notification also directs the State to prepare or update the CZMP by engaging reputed and experienced scientific institution(s) or the agencies and in consultation with the concerned stakeholders. Consequently, the Government of Kerala entrusted National Centre for Earth Science Studies (NCESS), Ministry of Earth Sciences, Thiruvananthapuram for the preparation of Coastal Zone Management Plan for the State of Kerala following the guidelines in CRZ Notification, 2019.

Accordingly, preparation of the draft CZMP in 1:25,000 scale map identifying and classifying the CRZ areas within the respective territories in accordance with the guidelines given in Annexure-IV to the CRZ Notification 2019 has been taken up by NCESS, which involve public consultation. The subsequent guidelines issued by the MoEF&CC based on Office Memorandum 12-1/2019-1A III dated 26-06-2020 is to facilitate the State Government in updation of the CZMPs. As per the new guidelines, The CZMP database (shapefiles etc.) prepared as per the CRZ Notification, 2011 which have been scrutinized by the Technical Scrutiny Committee, finalized by the National Centre for Sustainable Coastal Management (NCSCM) and approved by the MoEFCC, shall be used as the base for revision or updation of the CZMP, as per the provisions contained in the CRZ Notification, 2019. The guidelines brought out clarity in the case of the Data to be provided

by the States/UTs to the authorized agencies, CRZ buffers, Processing of Census data, CRZ Classifications, Public consultation of draft CZMP updated or revised based on CRZ Notification 2019, format for CZMP report, approval process of CZMP etc.

2. THE STATE OF KERALA

Kerala, the Gods own Country lies in the southwest corner of Peninsular India and positioned between 8°17'30"N and 12°47'40"N latitudes and 74°27'47"E and 77°37'12"E longitudes. It is bound by the Western Ghats Mountain ranges to the east and the Arabian Sea to the west. The Ghats run parallel to the west coast at a distance ranging from 40-80 km. Kerala is spread over a total area of 38,863 sq.km, having significant stretches of water bodies. Altitudes ranges from below sea level (the Kuttanad area) to 2,695 m and the terrain falls into three well marked divisions: (a) the high ranges of the Western Ghats in the east with undulating hilly tracts, marked by long spurs, extensive ravines and dense forest, (b) the midland occupies with plantations and cultivated plains intersected by numerous rivers and streams, and (c) the coastal belt with dense settlements, coconut plantations and rice fields (Soman, 2002). The total population of Kerala is 33,406,061 (as per 2011 census) with a density of 859 per sq km. The density of coastal urban population is 4,228 per sq. km., as compared to the average urban density of 2,097 in the state. The coastal rural population density is 1700, far above the state average rural population density of 603 (Geevan, 1996). The coastline length of Kerala is about 590 km. Kerala, despite its small land area with long coastline studded with world's best string of beaches. It is bestowed with a vast network of backwaters, lagoons, natural lakes, rivers and canals.

The wetlands of the state are categorized into two primary groups namely inland and coastal wetlands. The total area calculated as wetlands was 127930.07 ha, of which the inland wetlands cover approximately 34199.57 ha and the coastal wetlands estimated 93730.50 ha (MoEF, 1990). According to recent estimates by different agencies on wetland categories such as water spread area, aquatic vegetation and turbidity, it is around 1762 wetlands in the state. Moreover, 2592 wetlands smaller than 2.25 ha had been also identified. As a result, the total wetland area estimated was 160590 ha (Anon, 2010). CED, 2003 had suggested the major wetland classification system for Kerala based on different parameters like location, physical extend, depth, salinity, biodiversity etc

(Kokkal, 2008).

Kerala is rich with 44 rivers (41 west flowing and 3 east flowing) cut across Kerala with their numerous tributaries. The rivers either debouch into the Arabian Sea through inlets directly or drained to the sea through estuaries/lagoons (backwater). There are 48 inlets along the Kerala coast out of which 20 are permanent, whereas the remaining 28 are seasonal (remain open only during the monsoon period of June – September). The seasonal inlets mostly remain closed during the fair season due to the development of spit along the inlets due to deposition from longshore sediment transport. Seasonal inlets are normally cut open during monsoon for discharging storm- water accumulated from rainfall reducing the coastal inland from flooding risk. Reduction in the supply of riverine sediments might have affected the stability of the south-west coast in recent years.

The backwaters as a part of wetlands which running parallel to the coastline is a characteristic feature of the Kerala coast. It can be described as a body of brackish, marine or hypersaline water impounded by a sandy barrier and having an inlet connecting it with the open sea. Backwaters form an attractive and economically valuable and ecologically significant feature of Kerala. During monsoon, the backwaters overflow into the sea, discharging sizeable quantities of sediments, whereas in summer sea water flows into the backwater over considerable distances. The Kerala Public Works Department (Water Resources of Kerala, 1974) has identified 27 backwaters and 7 lagoons in Kerala. Kerala State has fourteen districts of which 9 districts has seacoast on its west.

There are 14 District, 152 Block Panchayats, 941 Grama Panchayats, 87 Municipalities and 6 Municipal Corporations. Apart from this, Kerala has one Cantonment (Kannur). The fourteen districts in the state are distribute over 75 Taluks consisting of a total of 1535 Villages (1664 is including the Group Villages). It is interesting to note that 9 districts (Kasaragod, Kannur, Kozhikode, Malappuram, Thrissur, Ernakulam, Alappuzha, Kollam and Thiruvananthapuram) out of the 14 have Lakshadweep Sea as their western boundaries and therefore come under the purview of CRZ covering considerable parts of coastline. Apart from the 9 districts, some parts of Kottayam district are also under the purview of CRZ since its western boundary is along the banks of tidal influenced Vembanad Lake.

2.1 Thrissur District

The word *TRICHUR* is the anglicised version of Thrissur which again is the abbreviated form of *Thrissivaperur* meaning the abode of Lord Siva. The district got its name from the headquarters town. It has recently shed its anglicized name and is now known as *Thrissur*. The town is built on an elevated ground, at the apex of which is the famous “VADAKKUMNATHAN” Temple. A place of great antiquity, Thrissur was also known as “VRISHABHADRI PURAM” and “TEN KAILASAM” in ancient days. Thrissur with its rich history, cultural heritage and archaeological wealth is called the cultural capital of Kerala. The present Thrissur district is a part of erstwhile Cochin state. With effect from the 1st July 1949 a new district named Thrissur was formed with 6 taluks of the erstwhile Cochin State viz. Talappilly, Chittoor, Thrissur, Mukundapuram, Cranganore and Cochin-Kanayannur and taluks of erstwhile Travancore areas, namely, Kunnathunad and Parur. Thrissur district is bordered by the districts of Palakkad and Malappuram to the north and the districts of Ernakulam and Idukki to the south. The Arabian Sea lies to the west and Western Ghats stretches towards the east. The district has a sea coast of about 54kms long and also has the presence of considerable stretch of lakes and backwaters. Thrissur district lies between latitudes 10°10'00"N and 10°46'00"N and longitudes 75°57'00"E and 76°54'00"E. Total area of Thrissur district is 3032km². In terms of area, Thrissur ranks 7th among the districts in the State. The location map of Thrissur district is given in Figure 1.

2.1(a) Administration

There are two systems of administrative set up in the State – Revenue and local self-government. Under the revenue system the district is divided into Revenue Division, Taluks and Villages. However, for local administration, the district is divided into Urban local bodies (Municipal Corporation and Municipal Councils) and rural local bodies comes under the hierarchy of District Panchayat consisting of Block Panchayats defined with geographically contiguous cluster of a few Grama Panchayats. For the implementation of development activities, Panchayats are grouped under Community Development Blocks. Therefore, all these units viz., Taluks, Villages, urban and rural local bodies have their own relevance and importance.

Thrissur district is one of the districts in the Central Kerala, which is having two Revenue

Divisions (Thrissur and Irinjalakkuda) consisting of 7 Taluks (Chavakkad, Kodungallur, Mukundapuram, Thalappally, Thrissur, Chalakkudy and Kunnamkulam) and 255 Villages. There are 29 Villages under Chavakkad taluk, 18 under Kodungallur, 29 under Mukundapuram, 45 under Thalappally, 74 under Thrissur, 29 under Kunnamkulam and 31 under Chalakkudy. Under the Local Self-Government System, the district is divided into 1 Municipal Corporation (Thrissur) 7 Municipal Councils (Chalakkudy, Chavakkad, Guruvayoor, Irinjalakkuda, Kodungallur, Kunnamkulam, and Wadakkancherry), 16 Development Blocks (Anthikad, Chalakkudy, Chavakkad, Cherpu, Chowannur, Irinjalakkuda, Kodakara, Mala, Mathilakam, Mullassery, Ollukkara, Pazhayanur, Puzhakkal, Thalikulam, Vellangallur and Wadakkanchery) and 86 Panchayats. The area encompassing the Thrissur city and surroundings forms the headquarters of the district.




Figure 1: Location map of Thrissur District

2.1(b) Physiography

As a continuum of the State, the district too can be divided into three distinct physiographical units such as the coastal plains, the midlands and the eastern highland regions. The coastal plains

form the low land areas adjacent to the Lakshadweep Sea.

Thrissur District consists of sub-micro regions, such as, (i) Thrissur Coast (ii) Thrissur Plain (iii) Wadakkancherry Upland and (iv) Kodassery Forested Hill and (iv) mountainous region.

(i) Thrissur Coast: This region lies along the coast of Thrissur district. It has its boundaries with Malappuram coast in the north, Thrissur plain in the east, Cochin coast in the south and the Lakshadweep Sea in the west. This coastal plain is drained by rivers like the Bharathapuzha, Wadakkanchery, Manali, Kurumali and Chalakudy, besides a network of canals and backwaters. This plain slopes very gently towards the west. The maximum height (55 m) is found in the central-east portion i.e., at Kodannur village of Thrissur taluk. The low-lying land, known as Koleland, lies as a continuous belt from the central portion towards the coast where the spot heights are 3 m, 10 m and 11 m etc. Paddy and coconut are the important vegetation in the region.

(ii) Thrissur Plain: This region is bounded by Pattambi undulating plain in the north, Wadakkancherry Uplands and Kodassery Forested Hills in the east, Periyar- Muvattupuzha Rolling Plain in the south and Thrissur Coast in the west. The region which is parallel to Thrissur coast has the maximum height of 118m at Amballur village of Mukundapuram taluk and has the minimum height of 43m at Muringur Thekkummuri village of Mukundapuram taluk. The central portion is an elevated area while the southern portion is low in height. The average height of this region is 75 m and sloping gently towards the west. This region is drained by the tributaries of the Bharathapuzha, Wadakkancherry, Manali, Kurumali and Chalakudy rivers. Paddy, coconut and cashew are the important vegetation of this region.

(iii) Wadakkancherry Uplands: This region has its boundaries with the Palakkad Gap on the north and the east, Kodassery Forested Hills in the south and Thrissur Plain in the west. It is not a high mountainous tract but only uplands having outlying hills in the eastern extreme of this region. Some knolls and hillocks are found here and there. The maximum height of this region is 208 m in Kondazhy village of Talappilly taluk and the minimum height is 52m in its southern portion. Vazham irrigation project is located on a lab-land in the southern portion of this region which is sloping on all sides. However, towards the south its slope is gentle and through this spur NH-47 runs connecting Thrissur with Palakkad. Areca nut, cashew, rubber, teak etc. are the main vegetation of

this region. Coconut and paddy are also cultivated here.

(iv) Kodasserry Forested Hills: This region lies in the south-eastern portion of the district forming its boundaries with Wadakkancherry Uplands in the north, Chittur Forested Hills and Tamil Nadu in the east, Malayattur Forested Hills and Anamalai Forested Hills in the south and Thrissur Plains in the west. This forested hill has wet semi-ever-green forest. In its northern and southern ends Peechi and Peringal kuthu reservoirs are located. This forest highland which is a part of the Western Ghats, forms the catchment areas of the Manali, Karuvannur and Chalakudy rivers. River terraces are seen in the middle portion. Numerous 'V' shaped valleys are located in this tract, which determines the course of various streams. Scarp slopes, waterfalls and rapids which structurally control and regulate the river directions are found mostly in the middle and southern portion of this region. The region in totality is highly erosion prone. The maximum height (1439 m) is found in the southern end of the reserve forest and the minimum height (416 m) is found in the northern end at the Attur village of Talappilly taluk. It slopes towards the north. Rubber and teak plantation are the vegetation cover of the area especially in the western end of this region.

(v) Mountainous region: The mountains of the district form an irregular outline along the Western Ghats with ridges and conical peaks. The Machad hills in Talappilly taluk, Paravattani hills in Thrissur taluk and Palappilly, Kodassery and Athirappally hills in Mukundapuram taluk are worth mentioning. There are about 30 peaks in the district with heights ranging between 330 metres and 1439 metres. Karimalagopuram, the highest peak in the district (1439 metres) lies at the boundary of Chittur taluk of Palakkad district and Mukundapuram taluk of Thrissur district.

2.1(c) Rivers and drainage characteristics

The important rivers in Thrissur district are Bharathapuzha, Keecheri, Puzhakkal, Karuvannur and Chalakkudy.

Bharathapuzha River, the second longest river of the State takes its origin at an elevation of +1964m above M.S.L from Anamalai hills and flows through the districts of Coimbatore, Palakkad, Malappuram, Thrissur and joins the Arabian Sea near the Ponnani town. Its main tributaries are the Gayathripuzha, Kannadi River or Chitturpuzha or Amaravathi, Kalpathipuzha and Thuthapuzha.

The Gayathripuzha has four main sub tributaries viz., Mangalam River in which Mangalam dam is located, Ayalurpuzha in which Pothundy dam is located, Vandazhipuzha, Meenkada River in which Meenkada dam is located and Chulliyar in which Chulliyar dam is located. Cheerakuzhi weir is located across the Gayathri River near Pazhayannur. The total length of the river is 209km with a catchment area of 6186 sq.km. About two-thirds of the drainage area of the basin i.e., 4400 sq.km lies in Kerala State and the balance 1786 sq.km. in Tamilnadu.

Keecheri River, which is also known as Wadakkanchery River on Alurpuzha is one of the smallest rivers in the State and is practically dry during summer. The river originates from Machad Malai at about +365m elevation in the upper reaches of Talappilly taluk forming part of the Western Ghats. The only important tributary of Keecheri River is Choondal thodu. The total length of the river is 51km. It has a total drainage area of 401sq.km.

Puzhakkal River which drains into the Kole lands of Thrissur district, is formed by the confluence of parathodu, Poomalathodu, Naduthodu and Kattachirathodu. Parathodu and Poomalathodu have their origin in the hills of Killannoor village at an elevation of +150m. Naduthodu rises from Manalithara hills on the South side of Machadmalai at an altitude of +525m while Kattachirathodu rises from below +75m near Mudikotty. It has a total length of 29 km and drainage area of 234 sq.km.

Karuvannur River originates from the Western Ghats and is fed by its two main tributaries namely Manali and Kurumali. The Manali originates from Vaniampara hills at an elevation of +365m. Chimony and Muply are two sub tributaries of the Kurumali originating from Pumalai at an elevation of +1100m. Karuvannur River has a length of 48km and drains an area of 1054 sq.km.

Chalakydy River has a length of 130km formed by the confluence of five streams namely Sholayur, Parambikulam, Kuriar (or Thekkadi), Anakkayam and Karappara, all of them originating from the Anamalai hills of Western Ghats. The river flows through the forest lands for the most part for a distance of about 80 km and then through fertile tracts for about 32 km. It then joins the right arm of Periyar River at Elanthikara about 10 km east of Kodungallur. The Hydro-electric projects at Peringalkuthu and Sholayur are the two main projects of the river. This river has a drainage area of about 1704 sq.km out of which, 1404 sq.km lies in Kerala State and the rest 300sq.km, in Tamilnadu.

2.1(d) Coastal Wetlands

Apart from the estuarine part of the rivers joining the Lakshadweep Sea, other important backwater network in the district is the Kole wetland system. The Kole wetland system of Thrissur district in Kerala, is an unique wetland system extending for an area of 11000ha in Thrissur and Malappuram districts, which is an important rice- growing tract of Kerala. The name Kole refers to the peculiar type of cultivation practice carried out on these lands. Kole in the regional language Malayalam indicates bumper yield or high returns in case floods did not damage the crop. Kole lands extend from the northern bank of Chalakudy River in the south to the southern bank of Bharathapuzha River in the north. A major portion of the area is flat, and it remains flooded during a period of six months from June to November. Four rivers namely Keecheri, Puzhakkal, Karuvannur and Chalakudy drain into this wetland system. These lands were formerly shallow lagoons, which gradually got silted up. The wetland is facing innumerable problems including flooding, salinity intrusion, deterioration of canals etc.

The area from Velukkara in the south on the Chalakudy riverbank in Mukundapuram Taluk to Mullassery of Chavakkad Taluk and Tholur-Kaiparama areas of Thrissur taluk is designated as ‘Thrissur kole’ and the contiguous area from Chavakkad and Choondal to Thavannur, covering Chavakkad and Thalapally taluks of Thrissur district and Ponnani Taluk of Malappuram district form the ‘Ponnani kole’. This area lies parallel to the sea and flood waters are mainly brought by two rivers namely Kechery and Karuvannur, which finally drains into the sea. A network of main and cross canals provides external drainage and connects the different regions of the Kole to the rivers. It is a basin with a saucer shape flanked by laterite hills in the western and eastern margins. Some portions of Kole area exhibit a lacustrine environment and contains black carbonaceous clay with many plant parts and in some places withered tree trunks. Also, the subsurface exploration of Kole land shows fine sandy deposits due to which mining has become an important activity in certain parts of the Kole lands recently. It is known that the soils in the Kole area, especially those in the flood plain are heavy clay in texture and acidic in reaction. The presence of organic peat layer in the sub-surface of the soil profile has made soil extremely acidic.

2.2. Demography and Socio-Economic Activities

Kerala has 33.38 million population as per the Census 2011. It is the 13th most populous State in India with an overall population density of 860 per square kilometer. The State of Kerala accounts for 2.8% of India's population but it contributes nearly 4% of the Indian economy. Kerala is occupied with three times more dense settlement than the rest of the country. The coastal regions are more populated than the mountains and eastern hills of the State with 2.5 times the overall population density. Kerala has a Human Development Index of 0.79, which is "very high" and the highest in India. Kerala also has the highest literacy rates among all Indian states at 98.9% and a life expectancy of 74 years which is among the highest in the country. Kerala has experienced a rapidly dropping rural poverty rate, which fell from 59% in the mid-1970s to 12% by 2010, while the overall poverty rate fell 47% between the 1970s and 2000s, compared to a drop of just 29% in total poverty in the country. As per 2011 census, Kerala is the most literate state in India having literacy rate of 93.91%.

While Kerala has what appears to be rapid growth by the standards of most areas, its 4.9% decadal population growth rate in 2011 was the lowest in India and less than one-third of the India average of 17.64%. Between 1951 and 1991, Kerala's population more than doubled from 15.6 million to 29.1 million, reaching 33.3 million by 2011. Kerala is currently heading for zero growth in its population, as the state has a meager fertility rate and a stabilizing death rate. In 2021, census figures predict Kerala might record negligible population growth, which will be a first in India. The population is advanced with literacy and educational attainment. The various sectors such as agriculture cash crop production, animal husbandry, aquaculture, fisheries, micro enterprises and large-scale industries, tourism etc. plays important role in the economy of Kerala. This state is unique in many respects among the states of India, one of which is its settlement pattern, characterized by a rural-urban continuum. Applying the "continuous method" to study spatial change in the occupational structure across rural, small towns and large urban units (comprising of cities/big/medium towns and agglomerations), it is interesting to note that economic diversification in general and manufacturing in particular, has been fairly rapid in rural areas.

As per the Census 2011 data, Thrissur district has a population of 31,21,200 persons, which stands the 4th place among the districts in the State. Thrissur also ranks 4th in terms of total households (759210) among the districts. In terms of population per sq.km, the district has recorded

the 7th position (1031) as against the State Density (860). The district has recorded the 9th rank in total (35.1 per cent) and 10th position in female work participation rate (18.7 per cent) and the 7th rank in male work participation rate (53.3 per cent). The marginal work participation rate of the district is 15.2 per cent. The district has the 2nd rank in the percentage of main workers 84.8 per cent. With 95.08 per cent, the district occupies the 7th position in literacy rate. The 6.7 percent of workers are agricultural labours, which occupying the 13th rank and the district occupies the 11th position in the percentage of cultivators (3.8 percent).

2.3. Coastal Geomorphology and Ecosystem

The shoreline of Kerala is generally straight, trending NNW-SSE, with minor variations. The various coastal geomorphological units are beaches, beach cliffs, stacks, islands, shore platforms, spits, bars, beach ridges, estuaries, lagoons and tidal flats. The beaches are mostly sandy and dynamic in nature. Throughout the coast exists narrow stretch of beach except in cliff areas. In areas like Kovalam, Vizhinjam, Varkala, Ezhimala, Bekal etc. the headland is directly abetting the sea where the wave break occurs along the foot of lateritic cliff. The height of the cliff may be of 20 m or more. In some coast having cliffs, there are numerous stacks protruding into the sea in nearshore as well as in offshore regions. These stacks are the vestiges left behind after an island or head land portion which has been eroded out or still receding. Around Mahe and Thalassery, these stacks are found aligned nearly parallel to the coast. In lateritic coast offshore islands are observed in certain locations. The Green Island located in the offshore of Thalassery coast is a similar type of island and have continuity with the mainland with a string of stacks. The constant wave attack on the neck portion resulted discontinuity of the stacks and becomes an island. The coastal wetlands, backwaters and estuaries along the coast of Kerala are rich, biologically and ecologically diverse as well as economically significant ecosystems which plays important role in livelihood of people. The major backwaters in Kerala are Vembanad, Ashtamudi, Kayamkulam, Akkulam, Kadinamkulam, Anchuthengu, Edava, Nadayara, Paravur, Vattakayal, Chettuva, and Valiyaparamba.

Thrissur district has in general an undulating topography with a regional westerly slope. Physiographically, the district is divisible into three zones from West to East as the coastal plain, followed by the midland region and then the hilly region. The coastal plain having an average width of 5km and height varying from 0-8m is a depositional landscape characterized by landforms of

marine, fluvial and fluvio-marine origin. A number of strand lines (Palaeobeach ridges) alternating with swales, aligned more or less parallel to the trend of present-day shore line can be seen in the area which could be representing successive still stand positions of the regressive episodes of the sea. The level of crests of the ridges shows an overall descending trend towards the sea suggesting progradation of an advancing coast. The midland region has an average width of 30km and has elevation from 20-300m. The terrain is characterized by laterite mesas and laterite interflaves and has a rolling or gently undulating landscape. The terrain has been dissected to give rise to narrow valley flats. Remnants of four former erosion surfaces are ranging in age from Late Tertiary to Early Quaternary has been reported from this area. Along the Eastern margin of the midland relatively high and resistant structural hill ranges can be seen. The hill ranges in the Eastern part are resultant of the structural cum denudational process in the past. This terrain comprises of rocks of Archaean age and has a thin soil cover. Along the Northeastern part of the district the topography is matured with a very gently sloping terrain. This probably represents a dissected pediment. Some of the highest peaks along the Eastern boundary are the 926m hill, 1160m Pappatapara hill and the 1439m high Karimalai.

2.4. Marine Fishery Resources

Fisheries sector is recognized as one of the important sectors contributing significantly to the nation's economy. It is not only recognized as a powerful income and employment generator as it stimulates growth of a number of subsidiary industries, but also is a source of cheap and nutritious food earning foreign exchange too. It provides livelihood to approximately 14.49 million people in our country. Kerala is one of the prominent maritime States in India and is blessed with most productive portion of Arabian Sea with a continental shelf area of 39,139 Sq km. Many commercially important finfishes and crustaceans form the high value fish species. The projected value of total fish production from Kerala comes to 706.882 MT by 2019-20. Over exploitation is one of the major threats in marine fishery resources. If the rich underexploited demersal and pelagic resources are utilized effectively, Kerala can brag the top position in total fish production and contribute fruitfully towards the economic development of the State. According to the available estimates of potential fishery resources of the West Coast, particularly in the south-west coasts, Kerala possesses the richest fishing grounds in the region. Marine fish landings for 2014-15 in Kerala was estimated at 524468 MT and that for 2013-14 was 522308 MT. Contribution of various fish resources include,

pelagic fin fishes 361956 69 MT, demersal fin fishes 49416 9 MT, crustaceans 65955 12 MT, molluscs 34057 6 MT and miscellaneous 13084 MT.

Kerala holds the second position in terms of fisherfolk population, among the nine maritime states in our country. A great deal of Kerala's economy depends on fishing for subsistence, livelihood and employment. Fish consumption in Kerala is four times the national average and the production share of Kerala is the second largest in the country with 16.6% of India's total marine exports. The total populace of fisherfolk residing in the state of Kerala is estimated to be 11.11 lakh, which includes 8.55 lakh in the marine sector and 2.55 lakh in the inland sector (GOK, 2015). Out of this, the number of active fishermen is 2.28 lakh (1.90 lakh in marine sector and 0.42 in the inland sector). Currently, there are 222 fishing villages in the marine sector and 113 in the inland sector, where fishing and its allied sectors provide livelihood to a vast majority of population (GOK, 2015). Primarily the fishers depend on fishing as the prime source of income. However, 12% of the fisherfolk generates additional income from allied fishing activities like marketing/repairing nets, fish vending/processing and other fishery related activities. The socio-economic condition of the fisherfolk in the state is sad, when compared to the general section of the population. Most of them are in the grip of subsistence economy and indebtedness due to socio-economic constraints, education and depletion of fishery resources. The density of population in the coastal area is 2168 persons per km².

Thrissur district has a long sea coast of about 54 km lying between Punnayurkulam in the north and Azhikode in the south. It is 9.2 percent of the total length of coastline in the State. Azhikkal is the only minor port on this coast. Chettuvai, a coastal village of the district has recently been developed into a fishing centre with facilities for mechanised fishing. Main fishing centres in the district are Chavakkad, Nattika and Kodungallur. Mechanised boats are also engaged for fishing in the district. Azhikode, Nattika, Vadanapally, Kadapuram, Chettuvai, Blangad and Puthankadapuram offer facilities for harbouring the mechanised boats. The district has 18 coastal fisheries villages and 8 inland fisheries villages. There is Shrimp Hatchery at Azhikode. As per the 2019-20 estimates of Fisheries Department, the fishermen population in the district is 77,274 with a breakup of 57,097 from marine and 20,177 from inland sector including the male, female and children. According to 2018-19 report, as per the Fisheries Handbook 2020, the annual fish production was 68170 MT from marine and 21589 MT from inland sector. There are 23 fish landing centres in the district and two

hygienic fish markets being managed by Mala Panchayat and Irinjalakuda Municipality. In the district, there are two Department hatcheries at Peechi in Pananchery Panchayat and a Regional Shrimp Hatchery at Azheekode in Kodungallur Municipality. ADAK is managing two fish farms in the district namely Government Fish Farm at Kadappuram in Thrissur Corporation and a Model Shrimp Farm & Training Centre at Poyya in Poyya Panchayat. MATSYAFED is managing a freshwater Prawn Hatchery at Kaipamangalam. Fishermen development welfare co-operative societies are functioning in this district extends all basic support and assistants for the development of inland fish farming.

2.5. Biodiversity of Thrissur district

Thrissur District has a 54 km long coastal belt, a few river mouths and unique plains with Kole lands which is the abode of natural aquatic and wetland species of flora and fauna. In the Kole wetland system of the district, the major formations of biodiversity are seen in the backwaters and associated marshes, ponds, streams, etc. Thrissur district with vast stretches of evergreen forest provides the natural habitat for a variety of plants and animals. Valuable medicinal plants, economically important trees, diverse group of mammals, birds and reptiles have the necessary ecological background to grow. There are much disturbed patches of mangroves also in the district with characteristic species of *Avicennia*, *Excoecaria*, *Kandlia* and *Carallia*. Terrestrial formations in the midlands are more characteristic of Agri biodiversity, apart from few elements of natural flora and fauna occurring in barren areas and gaps of cultivated lands. Many sacred groves in Thrissur district manifests past biodiversity. The natural forests in the district are represented by moist deciduous, semi evergreen, evergreen, savannahs and the southern wet temperate forests. As per the available literature the district has rich biodiversity showing plant diversity with 4000 species of flowering plants, 350 grass species, 15 bamboo species, 9 reeds species, 214 orchid species, 4 gymnosperms, 200 ferns and fern allies, 200 liverworts, 231 algae, 1044 fungi and 800 lichens. Among the animals, 48 species of large and medium sized mammals, 475 bird species, 101 aquatic birds, 60 reptiles Genera, 30 lizard (endemic) species, 57 snake (endemic) species, 87 amphibian (endemic) species, 84 freshwater fish (endemic) species and 313 butterflies.

2.6. Pollution and Waste Management issues

Kerala Solid Waste Management Project carried the waste quantification and characterization to represent the bulk waste generators of the State namely Household, Commercial and Institutional. The waste generation rate per capita in municipalities varies from 364 grams/capita to 456 grams/capita. Low waste generation is noticed in urban local bodies of highland areas. Highly urbanized Municipalities generate above 450 grams/capita and the City Corporation generates around 545 grams/capita. Domestic waste contributes 55-65percent of total waste, while commercial establishment and markets are the second-highest generators of waste. The average waste generation rate in Municipalities is 419 gm/capita/day whereas, the Municipal Corporation areas is 545 gm/capita/day. Based on the 3 broad categories of geographical regions (lowland, midland & highland), the waste generation rates are higher in the coastal belts, which is around 545 gm/capita/day in Municipal Corporation areas whereas, the waste generation rate in the midland belt is about 454 gm/capita/day and it is about 383 gm/capita/day in highland areas.

Mainly, the waste management includes the management of biodegradable waste, management of recyclable wastes, management of non-biodegradable and non - recyclable wastes and the management of biomedical wastes. As per the Solid Waste Management Rules, 2016, Centralized Windrow composting systems exists at Brahmapuram, Kochi and at Njaliyan parambu, Kozhikode. However, household level decentralized solid waste management facilities do exist at Kochi and Kozhikode Corportaion. More than 70% door to door collection of dry waste is achieved for households in 84 urban local bodies and for establishments in 70 urban local bodies as in May 2022. Haritha Karma Sena is working in 92 urban local bodies and 923 Grama panchayaths for collection of dry waste. For wet wastes disposal decentralized treatment methods such as aero bins, pipe compost, compost pits, kitchen bins, biogas plants etc. are followed. Dry wastes are collected, segregated and disposed through recyclers. In Kerala there are 147 plastic recycling units, 21 Steel mills, and 7 kraft paper units. Non-recyclable plastic waste is shredded in the Resource Recovery Facility and is used for the tarring of Public Works Department and Local Self Government Department roads. During the period 2016-2021, Clean Kerala Company Limited (CKCL) has produced 2399.13 T of shredded plastics and given to various agencies.

3. PURPOSE AND SCOPE OF CZMPS

The Coastal Zone Management Plans proposes a spatial planning framework for development by providing setbacks around sensitive eco-zones restricting development and other activities close to it. Setbacks require specific reference lines and boundaries for its meaningful implementation. The High Tide Line (HTL) forms the cardinal reference line for determining the setbacks for CRZ. The 50, 100, 200 and 500m CRZ lines landward from the HTL are the landward setback lines. In the case of inland Backwater islands and islands along the mainland coast, 20m from the HTL is uniformly demarcated. The Low Tide Line (LTL) and the Territorial water boundary (12 NM) form the setback lines towards the sea. The 50m line or width of the creek from the HTL has been demarcated along the tidal influenced water bodies that are connected to the sea and the distance up to which tidal effects are experienced, determined based on the salinity concentration of 5ppt. The CZMP has to be prepared in two scales (1:25,000 and 1:3960 or the nearest scale) in accordance with the guidelines given in Annexure-IV of CRZ notification 2019. The CZMP in 1:25000 scale with Survey of India Toposheets as base maps is required for formulating policy decisions. These are to be submitted to MoEFCC, Govt of India for approval after stakeholder/public consultations. The local level CZMP are to be prepared in 1:4000 with cadastral base maps and based on the approved CZMP. These local level CZM maps are for the use of local bodies and other agencies to facilitate the implementation of Coastal Zone Management Plans. The CZMP also has to incorporate the Hazard Line as demarcated by Survey of India (SoI) with a view to reduce the vulnerability of the coast. Critically Vulnerable Coastal Areas (CVCAs) demarcated by NCSCM is also incorporated into the CZMP prepared. **Shoreline of high, medium and low erosion stretches for such erosion prone areas will be added after receiving the data from NCSCM.**

4. COASTAL ZONE MANAGEMENT PLANS

The para 6 of the CRZ Notification 2019, numerates the following instructions for carrying out the CZMP of a State:

- (i) All coastal States and Union territory administrations shall revise or update their respective coastal zone management plan (CZMP) framed under CRZ Notification, 2011 number S.O.

19(E), dated 6th January, 2011, as per provisions of this notification and submit to the Ministry of Environment, Forest and Climate Change for approval at the earliest and all the project activities attracting the provisions of this notification shall be required to be appraised as per the updated CZMP under this notification and until and unless the CZMPs is so revised or updated, provisions of this notification shall not apply and the CZMP as per provisions of CRZ Notification, 2011 shall continue to be followed for appraisal and CRZ clearance to such projects.

- (ii) The CZMP may be prepared or updated by the coastal State Government or Union territory by engaging reputed and experienced scientific institution(s) or the agencies including the National Centre for Sustainable Coastal Management (hereinafter referred to as the NCSCM) of Ministry of Environment, Forest and Climate Change and in consultation with the concerned stakeholders.
- (iii) The coastal States and Union territories shall prepare draft CZMP in 1:25,000 scale map identifying and classifying the CRZ areas within the respective territories in accordance with the guidelines given in **Annexure-IV** to this notification, which involve public consultation. All developmental activities listed in this notification shall be regulated by the State Government, Union territory administration, local authorities or the concerned Coastal Zone Management Authority within the framework of such approved CZMP, as the case maybe, in accordance with provisions of this notification.
- (iv) The draft CZMP shall be submitted by the State Government or Union territory to the concerned Coastal Zone Management Authority for appraisal, including appropriate consultations and recommendations in accordance with the procedure(s) laid down in the Environment (Protection) Act, 1986 (29 of 1986).
- (v) The Ministry of Environment, Forest and Climate Change shall thereafter consider and approve the respective CZMP of concerned State Governments or Union territory administrations.
- (vi) The CZMP shall not normally be revised before a period of five years after which, the concerned State Government or the Union territory may consider undertaking a revision.

4.1. Demarcation of High Tide Line (HTL) and Low Tide Line (LTL)

The highest level horizontal positional and spatial accuracy in mapping and presenting the

HTL becomes necessary for field uses by CRZ implementing agencies. The agencies are looking for a planimetric accuracy approaching zero error. The different approaches now practiced in the country to demarcate the HTL are Tide level projection, using morphological signatures observed in the field as well as from the high-resolution satellite imageries. NCESS follows the approach as per the guidelines mentioned in the Annexure IV of CRZ Notification 2019. As per the Amendment to the CRZ Notification 2019: gazette notification no. S.O. 1422(e) dated 1st may, 2020 & no. S.O. 4886(e) dated 26th November 2021, In case there exists a bund or a sluice gate constructed in the past, prior to the date of notification issued vide S.O. 114(E) dated 19th February 1991, the HTL shall be restricted up to the line long along the bund or the sluice gate and in such a case, area under mangroves arising due to saline water ingress beyond the bund or sluice gate shall be classified as CRZ-IA irrespective of the extent of the area beyond the bund or sluice gate. Such areas under mangroves shall be protected and shall not be diverted for any developmental activities. The coastal morphological signatures are collected by field work as well as from the satellite imageries for the purpose of demarcation of HTL.

Morphological signatures are good indicators of shoreline oscillation and inundation of coastal waters, which could be used for identifying the HTL. The inundation of coastal waters on to the land and seasonal shoreline oscillations are dependent on coastal morphology. Shoreline remains stable and would not retreat significantly along cliffy coasts. The shoreline retreats up to the cliff base along pocket beaches. Artificial morphologies like seawalls confine the oscillation of shoreline along the line of the structure itself. Sandy beaches are prone to seasonal and long-term shoreline oscillation. Long term stability of the beach and the position of the stable part of the beach would be evident from morphological signatures such as berm and berm crest. This could be done by field methods and using combination of spatial data sources including satellite data. The HTL must be fixed with respect to certain reference points on the land. These reference points at sufficiently close intervals (preferably at least 1km along shore) have to be marked with respect to latitude-longitude and known points in the base map. Geomorphologic features like berm crest, cliff, headland, line of permanent vegetation, etc. are indicators of the reach of sea water into the land. Stable coastal protective structures like seawall also limit the intrusion of seawater. Hence High Tide Line (line of maximum reach of seawater into the land during spring tide) can be fixed in the field, with respect to these features and tied to the reference points, as detailed below:

a) *Landward (monsoonal) berm crest for beaches*

In all the well-formed wide beaches, one or more berms (which are nearly horizontal part of the beach developed through the deposition of sand by wave action) are usually observed. The seaward end of the berm at which a sudden downward slope is observed is termed a berm crest. When there is only one berm, it normally gets eroded during the monsoon with a berm crest on the landward side. But when there are two berms the landward berm is the monsoonal berm, which normally do not get eroded. Or else we can say that the erosion reaches only to the second berm crest. Since the tidal waters do not reach the coast beyond this landward berm crest, it is taken as the HTL. The distance to this point from the reference point is measured using the beach profile to fix the position of the HTL.

b) *Seawall/revetments/embankments*

In highly erosion-prone areas, no second berm is observed landward. Such locations will be protected mostly by seawalls. During monsoon season majority of these places are devoid of beaches. The waves impinge upon the seawall during the monsoon season, especially during the high tide. Thus, they are the artificial barriers stopping the waves/tides at the coast. Since the seaward part of the seawall in most cases is defaced due to erosion, the landward toe is taken as the HTL boundary in such locations. There are some locations with two or three lines of seawall, particularly in the accreting areas. The seaward seawall is considered here for the purpose. On the other extreme, in the case of continuously eroding sites there are lines of sea wall which are now in the sea. In such cases the landward seawall is taken. In order to facilitate the demarcation of HTL at seawall locations, the latter has to be clearly marked in the beach profile during coastal surveys.

c) *Permanent Vegetation Line*

Permanent vegetation develops on the stable part of the beach. There are several locations along Kerala coast, which has only one berm and the beaches undergo severe erosion during the monsoon, and yet not protected by seawalls. In such cases, permanent vegetation, particularly well grown coconut trees, which are the main vegetation species prevalent all along the coast, is used as an indicator. The part of the beach landward of monsoon berm crest, which is mostly stable, and the

line of permanent vegetation normally follows the line of monsoon berm crest which is the HTL.

d) *Coastal sand dune / paleo-aeolian dune*

Sand dunes are mounds, hills or ridges of sand that lies behind the part of the beach affected by tides. They are formed over many years when windblown sand is trapped by beach vegetation or other stationary objects. Sand dunes are habitat for coastal plants and animals. The size and morphology of coastal dunes is dependent on the complex interaction between controlling winds, sediment supply, and the geomorphology of the nearshore and beach environment. Mostly, dunes can be divided into those that form from the direct supply of sediment from the beach face (primary dunes), and those that form from the subsequent modification of primary dunes (secondary dunes). Sand dunes provides and storage and supply for adjacent beaches. They also protect inland areas from storm surges, hurricanes, floodwater, and wind and wave action that can damage property. Sand dunes support an array of organisms by providing nesting habitat for coastal bird species including migratory birds. The main secondary dunes include blowouts, parabolic dunes, and transgressive dune fields.

In Kerala, coastal inland areas have remnants of coast-parallel sand ridges manifesting the Holocene transgressive still stands of sea. North and Central Kerala coasts had such dispositions of strandlines of alternating ridges with swales. However, due to the demand of dense population in the coastal region and development activities, we could rarely see such raised dunes/ridges currently in Kerala except along the Pallikkara-Kanhangad coastal belt in the Kasaragod District. Another interesting feature witnessed is the foredunes bordering the beaches along most part of Kerala coast. They are seen in the backshore of the beaches as shadow dunes continuously being formed due to the sand blown out and trapped around any obstruction such as shrubs or grasses in the backshore.

e) *Mangroves*

Mangroves are unique plant communities comprising of evergreen trees and shrubs belonging to several unrelated families observed in tropical to subtropical intertidal regions, where constant tidal water exchange takes place. Mangrove ecosystem ecologically functions as a complex ecotone or interface zone between the terrestrial and marine ecosystems, exemplifying diverse

habitats, including microhabitats, characteristic of terrestrial, intertidal and aquatic environs. As an invaluable ecological system and for reasons of its rich biodiversity, economic and social standing for sustenance and survival of community people, sustainability of sea food, and shore-line stability, conservation of mangroves is of paramount importance. They exhibit remarkable adaptation for salt tolerance with a spread of around 1 lakh sq.km world over distributed in about 30 countries. Mangroves in India account for about 5 percent of the world's mangrove vegetation and are spread over an area of about 4,800 sq.km along the coastal States/UTs of the country. The best development of mangroves in India is along the east coast with nearly 57% (~2750 sq.km) of the mangrove ecosystem of the country. Along the west coast of the country occur 23% (~1100sq.km) of the Indian mangroves and the remaining 20% is around the Andaman and Nicobar Islands (India). The east coast, unlike the west coast, is endowed with the largest mangrove wetlands developed on larger river deltas created by the major east flowing rivers of the country.

Kerala once in the 1950's was blessed with a large spread of about 700 sq.km mangroves but has been declined considerably to around 20 sq.km. All along the Kerala coast there are a good number of small mangroves stands, though mostly in isolated patches, fringing the estuaries and backwaters (kayals); and around islets or along river margins in the coastline stretches. Kerala with its very limited extent of mangroves is in no way free from the current trends of degradation of mangrove systems in the country. Mangrove systems in Kerala exhibited a higher grade of heterogeneity in their environmental settings and ecosystem features. Mangrove systems are one of the most threatened habitats in Kerala, as anywhere else in the country, or in the world. There is confusion about the actual/exact extent of mangrove distribution in Kerala in the absence of a precise estimate of it.

There are 15 true mangrove species and 49 mangrove associates observed in the coastal brackish water areas of Kerala. The 15 true mangrove species belonged to 9 genera spread over 7 families. The family, Rhizophoraceae is the most represented one with 6 species belonging to 3 genera. Mangrove associates are generally observed in the fringe areas where the wetland nature is devoid of any salinity. Species like *Acanthus ilicifolicus*, *Excoecariaagallocha*, *Aegicerascorniculatum*, *Rhizophora mucronata*, *Sonneratiaapetalae* and *Acrostichumaureum* were found in all the districts of Kerala, whereas *Rhizophora apiculata* is widely distributed in Kannur and Kollam districts but not found in Malappuram. *Avicennia officinalis* is one of the

common species noticed in all the districts, however, this is not the case with *A. marina* which was not seen in Trivandrum and is one of the threatened mangrove species in Kerala. Out of four species belonging to the genus *Bruguiera*, *B. cylindrica* has relatively wide distribution, however, it is not recorded from in Kottayam district. *B. parviflora* has wide distribution in the northern parts of Kerala which is not present in Trivandrum, Kollam, Alappuzha and Kottayam. *Kandeliakandalis* also a rare species which is distributed in all districts except Trivandrum, so also *Sonneratiacaseolaris* which is found in five districts namely Trivandrum, Kollam, Alappuzha, Kannur and Kasargode, whereas *S. alba* is becoming endangered due to its small populations in the districts of Ernakulam, Kozhikode, Kannur and Kasargode. *Lumnitzeraracemosais* one of the rarest mangrove species in Kerala found in four districts namely, Trivandrum, Kollam, Alappuzha and Kannur.

Sometimes, small creeks or pockets or far inland areas in the upriver vicinities or close to coastal waterways or canal works, harbour mangrove systems, which may apparently remain dry for considerable time, but remain intact-unless it is wantonly degraded by man-since the water table of area or site is generally just below surface. Both the fringe and the riverine mangroves have significant productivity status as they turn out high amount of organic matter. The wetland bays or pockets, of small or large areas, associated with them are affected by freshwater runoff from adjacent upland areas and rivers, as well as sediments and nutrients transported by the river inflows.

The mangroves and the mangrove wetland system in and around Kadalundy, offer congenial habitats or home grounds for many and varied faunal communities, which remain well integrated in a natural web of food chains, right from the detritus feeders and primary consumers (herbivores) to secondary, or tertiary consumers (carnivores). The mangroves and the wetland system being located in, somewhat like, an isolated setting, comparatively much away from the human settlement areas, the ecosystem premises are amply used as the homing environs by a wide variety of animals, both invertebrates and vertebrates. Kozhikode district has a better distribution of mangrove stocks and is third in position with considerable areal extent. The stands are in varied configuration owing to the impacts of degradation. The wanton degradation of mangrove systems in this district portrays a picture that evokes the careless attitude and disregard for mangrove wealth. The mangrove depletion at many sites in the district has gone to the extent that the functional ecosystem role of those stands has been narrowing down, over the years, with its tell-tale implications on hydrological and biotic features. A good part of the mangrove systems is thriving very much in the developmental limits of

the urban settings of the district; the existing mangrove patches within these areas are steadily and systematically being removed in the guise of developmental activities of the district. The important mangrove growing areas in the district include the mangroves of the Beypore and Kallayi River sides, mangroves at Eranhipalam-Kottooly areas close to the Canolly Canal, and the mangroves associated with the Kolavipalam-Kottapuzha estuary and backwater embayment. Mangroves of Chaliyar and Kallai River sides/estuaries in which the upper reaches of river stretches in continuum with the estuaries of the Chaliyar River (Beypore estuary) and the Kallayi River have disrupted stretches of mangroves along their margins. The estuarine premises of these rivers are almost entirely devoid of mangroves, owing to varied kinds of pressures from urban settlements, port developmental and other industrial activities.

Species like *Acanthus ilicifolius*, *Excoecariaagallocha*, *Aegicerascorniculatum*, *Rhizophora mucronata*, *Sonneratiaapetalae* and *Acrostichumaureum* are found in the district, which are available in all the districts of Kerala. Azhikode is the extreme south region of Thrissur district, in the nearby area, patches of mangroves were seen in Karupadanna, Kodungallur, etc. *Avicennia officinalis* was common in all these areas. Poyya backwaters area still sustains some mangroves. Kadungachira is another mangrove area with presence of *Avicennia officinalis*, *Excoecariaagallocha* and *Acrostichumaureum*. Krishnankotta was a mangrove area where along with Karupadanna and Pullut (near Kodungallur) small patches of mangroves are observed. Chettuva is an important area for mangroves in Thrissur district, where the estuarine environment supports mangroves which comes under Orumanayur Grama Panchayat of Chavakkad Taluk. The confluence of Chettuvapuzha and Karanjirapuzha with the Lakshadweep Sea, a significant stretch of mangrove vegetation has been recorded.

From Chettuva, species such as *Acanthus ilicifolius*, *Aegicerascorniculatum*, *Avicennia officinalis*, *Bruguiera cylindrica* and *Rhizophora mucronota* were observed, whereas abundance of *Avicennia officinalis*, *Acanthus ilicifolius*, and *Acrostichumaureum* were reported from Azhikode. Mangroves of Pullutare rich in *Avicennia officinalis*, *Acanthus ilicifolius* and *Acrostichumaureum*. Poyya backwaters are abundant in *Avicennia*, *Aegiceras*, *Excoecariaagallocha* and *Acrostichum*. Chakkamkandam and Thekkan Palayur have *Avicennia*, *Acanthus* and *Acrostichum*. Venmanad area are with the formations of *Acanthus*, *Avicennia*, *Rhizophora* and *Acrostichum*. Near Enamakal Regulator, a small patch of mangrove with presence of *Rhizophora*, *Acanthus* and *Acrostichum* has

been found.

f) *Rocks, Headlands, Cliffs*

The 590km long Kerala coast is dotted in between by rocky promontories, headlands and steeply sloping cliffs. Except the coastal districts of Ernakulam and Alappuzha, all the other 7 districts have distribution of such rocks/headlands/cliffs to varied extent. At the rock outcrops, headlands and cliffs the water is quite deep that there is virtually no spatial displacement in the waterline. Hence, the High-Water Line available in the topographical maps (transferred to the base map) can be taken as such. However, at the eroding laterite cliffs (e.g., Varkala, Paravoor, Thalassery in Kerala), the latest position of the toe is taken from the cross section measured at the respective sites. This is to be verified against the satellite imagery and transferred to the base map.

The cliffs and rocky promontories are not present along the Thrissur coast.

g) *Influence of Tidal action*

CRZ shall apply to the land area between HTL to 50 meters or width of the creek, whichever is less on the landward side along the tidal influenced water bodies that are connected to the sea. The tidal influenced water bodies as per the CRZ Notification 2019 means the water bodies influenced by tidal effects from sea in the bays, estuaries, rivers, creeks, backwaters, lagoons, ponds that are connected to the sea. The distance up to which CRZ is applicable shall be governed by the distance up to which the tidal effects are experienced which shall be determined based on salinity concentration of five parts per thousand (ppt) measured during the driest period of the year and distance up to which tidal effects are experienced. As per the Office Memorandum dated 26th June, 2019 of the MoEFCC, guidelines were issued facilitating updation of CZMPs, according to which, the CZMP database prepared as per the CRZ Notification 2011 shall be used as the base for revision or updation of the CZMP, as per the provisions contained in the CRZ Notification, 2019. Therefore, the tidal effects as marked in the CZMP prepared as per the CRZ Notification 2011 forms the basis for the updation of the CZMP currently being prepared. The tidal limit of various tide influenced waterbodies of the Thrissur district is given in the table below in terms of location with latitude and longitude.

Table: Salinity/CRZ limit along the inland water bodies in Thrissur District

Sl. no	Name of Waterbody	Latitude	Longitude
1	Chetwai Puzha	10°32'19.76"N	76° 4'5.54"E
2	Chetwai Puzha (tributary)	10°35'12.18"N	76° 2'8.16"E
3	Puzhakkal River	10°30'13.74"N	76° 6'30.59"E
4	Karuvannur River	10°23'49.38"N	76° 9'0.36"E
5	Muriyad-Moorkanal canal	10°20'19.59"N	76° 9'25.38"E
6	Shanmugham Canal	10°19'14.28"N	76°10'4.48"E
7	Pullut River	10°15'27.29"N	76°15'22.04"E

4.2. Demarcation of Ecologically Sensitive Areas

Guidelines for preparation of the CZMP specifies that the CZM maps shall clearly demarcate the land use plan of the area and map out the Ecologically Sensitive Areas (ESAs) or the CRZ-IA areas as per mapping made available by NCSCM to coastal State and Union territories. Coastal ecosystems provide a variety of ecosystem services for humans; however, these systems are susceptible to both terrestrial and marine factors because they are situated in the coastal ecotone. Consequently, coastal marine ecosystems are very sensitive to environmental change and human activities. Constructions for coastal development are still often located in sensitive biological and ecological areas without much consideration of their impact. In this context, the CRZ Notification 2019 provides provisions to achieve coordinated development of the population, economy, and environment in the coastal area. Ecological sensitivity refers to the degree of reflection of ecosystem interference in human activities and changes to the natural environment; that is, the degree to which the ecosystem responds to environmental changes caused by the combination of internal and external factors. Through identifying such sensitive areas, conservation and management strategies could be developed that facilitate the sustainable use of coastal resources.

The different ecologically sensitive areas as listed in the CRZ Notification 2019 have been

assimilated from the previously approved CZMP maps as directed through the guidelines and further the changes occurred till recently has been captured using high resolution satellite data. The same has been field verified for accuracy assessment wherever required.

5. ECOLOGICALLY SENSITIVE AREAS/COASTAL LANDUSE

5.1. Mangroves

Mangrove stands of the Thrissur district exhibit only sublevel growth as degrading factors have always been limiting the processes of succession maintaining them in the disturbed condition. As per literature *Acanthus ilicifolius* and *Avicennia officinalis* were common in almost all mangrove areas in Thrissur district. *Aegicerascorniculatum* was rare and reported from Chettuva and Poyya backwaters. *Bruguiera cylindrica* and *Rhizophora mucronata* were present only in Chettuva estuary. In Enamakkal *Rhizophora* was rarely seen. *Excoecariaagallocha* was present in Kadungachira, Poyya and Krishankotta. *Acrostichumaureum* was present in almost all study areas except Chettuva. It was reported that among mangrove associates, *Derris uliginosa* was present in almost all areas in Thrissur district where mangrove occur. *Clerodendroninerve* was present in Chettuva, Poyya and Azhikode. *Thespesia populnea* was present in Azhikode, Chettuwei and Krishnankotta. *Ipomea biloba* was present in Krishnankotta and Azhikode. *Sphaeranthus indicus* was common in Poyya and Krishankotta.

The total extent of mangroves in Thrissur district is 1.315048 Km², which is a moderate presence among the coastal districts in Kerala. The distribution of mangroves is highly skewed with the southern part of the Thrissur District (Kodungallur, Puthenchira and Poyya) having about 60% of total mangrove presence. Kodungallur Municipality tops the list with 0.402508 Km² of mangroves, followed by the Puthenchira Panchayat (0.266299 Km²) and Poyya (0.164177 Km²). The central part of the Thrissur is almost devoid of mangroves. Moving towards north, the Panchayats such as Pavaratty, Engandiyur and Venkidangu and the Municipalities such as Chavakkad and Guruvayoor have sporadic presence of mangroves in the range of 0.028361 to 0.4055154Km². (Detailed table is annexed in Annexure 2).

5.2. Coral Reefs

Coral Reefs locations have not been reported from the Thrissur coast.

5.3. Reserve Forests

Reserve Forests have not been reported from the Thrissur coast.

5.4. Sand Dunes

Sand dune locations have not been reported from the Thrissur coast.

5.5. Salt marsh

Salt marsh locations have not been reported from the Thrissur coast.

5.6. Nesting Ground of Birds

The nesting ground of bird's have not been reported from the Thrissur coast.

5.7. Archaeologically important and Heritage Sites

An archaeological site is a place (or group of physical sites) in which evidence of past activity is preserved (either prehistoric or historic or contemporary). Archaeological sites are open museum for living history. Heritage is a broad concept that includes the natural as well as the cultural environment. It encompasses landscapes, historic places, sites and built environments, as well as biodiversity, collections, past and continuing cultural practices, knowledge and living experiences. The prime concern of Conservation, Preservation and Maintenance of ancient monuments and archaeological sites along the coastal region remains one of the objectives of the CRZ Notification.

There are four archeological or heritage sites in the CRZ area of the Thrissur district, viz., Chettuva Kotta, Kodungallur Kovilakam, Kizhthali Siva Temple and Kottapuram Fort (Given in the map – Annexure-I).

Chettuva or Manappuram Kotta is located 11 km, north-west of Thrissur. In 1714 the Dutch

took possession of the territory and thereafter its control changed several times between the Zamorin and the Dutch. The Dutch erected a fort of Chettuvai and named it 'Fort William'. The island and the fort were the scene of several fierce encounters between the Dutch and the Mysore rulers (Haider Ali and Tippu Sultan) for about 50 years till Colonel Martley finally cleared Chettuvai of the Mysorians in 1790.

Kodungallur Kovilakam is a palace of the royal family of the late [medieval] Kingdom of Kodungallur (Cranganore), in the modern-day Indian state of Kerala. Kodungallur was a feudal principality subordinate to the rulers of the Kingdom of Cochin from the latter half of the eighteenth century until Indian independence. The Kingdom of Kodungallur was under the protection of the Dutch government after 1707 for a few years before returning to its allegiance to the Zamorin. The Kodungallur Royal Family had two branches, at Chirakkal Kovilakam and at Puthen Kovilakam

According to Keralolpathi, Kizhthali Siva Temple existed during the Perumal reign (B.C 113-AD 343). During the time of the Chera dynasty, who were Shaivates, believers of Lord Shiva, the kings had many advisors, usually nambuthiris. These ministers or advisors customarily gathered at a Shiva temple called Thali. There were so many Shiva temples in and around this main temple such as Melthali, Nedyathali and Chingapuram Thali. Kizhthali temple was first destroyed by the Portuguese and then Dutch. Then Tipu's army destroyed it further, except the *garbagriham*, which still stands. In its original glory, the temple boasted of a Koothu Parambu, a Kalari Parambu and a Kalapura Parambu. Presently, it is maintained by the Department of Archaeology, while the religious functions are carried out by a private trust. The temple is situated on the Paravur-Kodungallur Road.

Kottappuram Fort, popularly known as Crangannoor (Kodungallur) Fort built by the Portuguese in 1523 was later captured and destroyed by the Dutch in 1663. It was located in a strategic position, on the mouth of the river Periyar before it joins the Arabian Sea. Hence, it was easy to control the ships and boats that pass through the river to the interior of Malabar and vice versa, from this fort itself. Kottappuram Fort played a significant role in many wars between the Zamorin and the rulers of Kochi. In 1662 the Dutch fleet had made an attempt to conquer this fort from the Portuguese and they defeated the Portuguese in 1663. In that heavy fight, the Kottappuram fort got damaged. After taking over the fort, the Dutch reduced it to the minimum and used it as an outhouse to guard their trade ships. The remnants of the original fort envisage that it was 18 feet in

thickness and made of laterite. When the interest of the rulers of Mysore turned towards Malabar, Hyder Ali negotiated with the Dutch for the purchase of the forts at Kottappuram and one at Pallippuram. When Tippu Sultan came to be in possession of the Malabar coast it became imperative on the part of the Travancore Government to possess these forts in order to protect the country from the invasion of the Mysore rulers. Then Travancore King Ramavarma (Dharmaraja) (1758-1798) purchased these two forts on 31 July 1789. The agreement was executed by Raja Kesava Dasa, the Diwan of Travancore and John Gerard Van Angelbeck, the Dutch Governor. In 1909, the Department of Archaeology of Travancore erected a memorial pillar inside the fort and decided to preserve the fort as public property as it was in a ruined state at that time.

Altogether, the area under the Archaeological/Heritage site in Thrissur district is 0.011031 Km² (Detailed table is annexed in Annexure 2).

5.8. Seagrass

Seagrass locations have not been reported from the Thrissur coast.

5.9. Mud flats

Mud flats locations have not been reported from the Thrissur coast.

5.10. Turtle Nesting Grounds

Two beach stretches in Orumanayur Panchayat and one small beach stretch in northern part of Chavakkad Municipality beach has been identified as turtle nesting site along the Thrissur district (locations provided in the map given in Annexure-I). Local information gathered confirms that turtles do not appear regularly during nesting period of November and December every year. The total area of turtle nesting for the three stretches accounts for 0.043373 Km². (Detailed table is annexed in Annexure 2).

5.11. Inter-Tidal Zone

Tides play very important role in determining the biodiversity of and fertility of coastal and estuarine ecosystems. Intertidal zone is the area between the high tide (HTL) and low tide lines

(LTL) as per the CRZ Notification. Intertidal zone exists wherever the tidal effects are experienced. The intertidal zone is an ecosystem where a multitude of organisms living on the shore/banks survive changes between high and low tides. The tidal ranges are low in the southern side of the west coast of India and as we move northward, its amplitude increases. At Kochi, the ranges are of the order of 1m. The tidal range increases northward and reaches to more than 2m at Marmagao. At Mumbai, maximum ranges in tidal elevations are of about 5 m. Kerala coast being microtidal in nature with tidal amplitude around 1m with slight increase from south to north, the extent of intertidal area by the sea and inland water bodies are limited.

In the case of Thrissur district, intertidal area within the CRZ-IB category is bifurcated into the intertidal areas by the sea coast as well as the tidal waterbodies which accounts for 11.82725 Km². Apart from this, certain stretches of Pokkali fields are available in the Thrissur district accounting for an area of 0.644082 Km² (provided in the Table in the Annexure-2), where alternating seasonal practices of aquaculture and paddy cultivation is being done. Puthenchira Panchayat tops the list with 3.87726 Km² spread of intertidal area, followed by KodungallurMunicipality with 2.087885Km². Except Thanniam, Anthikkad and Kattoor panchayats, all other local bodies have varying spreads of intertidal areas. In the case of Pokkali category of CRZ IB, Poyya panchayat (0.486392 Km²) followed by Puthenchira panchayat (0.089757 Km²) have considerable spread. (Detailed table is annexed in Annexure 2).

5.12. Salt pan / Aquaculture ponds

Though aquaculture ponds are available in Thrissur district, they are mostly classified under the intertidal areas (CRZ IB) or as CRZ IVB due to their alternating use for aquaculture and agriculture. The same sort of categorization adopted in the approved CZMP of 2011 is being followed in the updation process.

6. METHODOLOGY FOR PREPARATION OF CZMP

As outlined in the Annexure-IV, preparation of the CZMP has been undertaken in 1:25000 scale using the base grids of the Survey of India (SOI) topographic sheets. Wherever 1:25000 scaled toposheets are not available, the 1:50000 SOI toposheets were enlarged accordingly to compose the base maps. The base maps were georeferenced as per the datum and projections specified in the

guidelines. The cadastral maps of the villages (1:3960 or nearest scales as per availability) likely to be within the purview of CRZ have been appropriately georeferenced to maintain the horizontal accuracy required. To minimize the RMS error during the georeferencing, maximum number of control points were obtained from the field using GPS (combination of methods using long static DGPS, short static DGPS and RTK obtaining acceptable precision resolving ambiguities in the post-processing techniques) to define the location in terms of latitude and longitude geodetic points in DMS format with second decimal accuracy in seconds (X & Y as cartesian coordinates with submeter accuracy). Wherever, disparity has been noticed in the hardcopy scanned cadastral image, georeferencing has been done by seeding maximum control points adjusting with the physical signatures discernible on the high-resolution satellite image which has been used as a reference image.

6.1. Field mapping and map preparation

The field mapping has been performed with hard copy of the georeferenced cadastral sheets to match the mapping scale with the ground space distance. Hard copy of the satellite images to a matchable scale has also been printed out to refer simultaneously during the field survey which comes handy in matching with the co-locatable ground features. The field surveying becomes confident with the combination of both cadastral and satellite images as well as with tying up the coordinates obtained from the GPS. All the essential features and lines are captured using the GPS by tagging attributes while carrying out the field work so that it becomes easy collate the information into maps without any confusion or missing. The guidelines issued subsequent to the notification specifically clarifies that HTL, LTL, ESAs and Critically Vulnerable Coastal Areas (CVCAs) demarcated by the NCSCM, Chennai, and the Hazard Line as demarcated by the SOI, shall be used in preparation/updation of the CZMPs as required under the provisions of the CRZ Notification, 2019.

Timeline satellite images were verified for any considerable change in the HTL/LTL/ESAs. Significant changes in the terms of the reduction or increase of mangroves have been noticed at several places in the State, which has been verified in the field intensively. Based on the ground condition, the variations in the extent of mangroves as well as the changes in HTL/LTL has been marked using the GPS tracking *in situ*. Apart from digesting the changes in the CZMP, separate table

has been created to mark the changes at each location for scrutinizing the same at the vetting stage.

7. CRZ CLASSIFICATION

The CRZ Notification 2019 has classified the CRZ area in the following manner for the purpose of conserving and protecting the coastal areas and marine waters.

7.1. CRZ-1

CRZ-1 areas are environmentally most critical and are further classified as under:

7.1.1. CRZ-1 A

CRZ-1 A shall constitute the following ecologically sensitive areas (ESAs) and the geomorphological features which play a role in maintaining the integrity of the coast viz.:

- (i) Mangroves (in case mangrove area is more than 1000 square meters, a buffer of 50 meters along the mangroves shall be provided and such area shall also constitute CRZ–IA).
- (ii) Corals and coral reefs.
- (iii) Sand dunes.
- (iv) Biologically active mudflats.
- (v) National parks, marine parks, sanctuaries, reserve forests, wildlife habitats and other protected areas under the provisions of Wildlife (Protection) Act, 1972 (53 of 1972), Forest (Conservation) Act, 1980 (69 of 1980) or Environment (Protection) Act, 1986 (29 of 1986), including Biosphere Reserves.
- (vi) Salt marshes.
- (vii) Turtle nesting grounds.
- (viii) Horseshoe crabs' habitats.
- (ix) Sea grass beds.
- (x) Nesting grounds of birds.
- (xi) Areas or structures of archaeological importance and heritage sites.

7.1.2. CRZ-I B

The intertidal zone i.e., the area between the Low Tide Line and High Tide Line constitutes the CRZ-I B.

7.2. CRZ-II

CRZ-II constitutes the developed land areas up to or close to the shoreline, within the existing municipal limits or in other existing legally designated urban areas, which are substantially built-up with a ratio of built-up plots to that of total plots being more than 50 per cent and have been provided with drainage and approach roads and other infrastructural facilities, such as water supply, sewerage mains, etc.

7.3. CRZ-III

Land areas that are relatively undisturbed (viz. rural areas, etc.) and those which do not fall under CRZ-II, shall constitute CRZ-III and CRZ-III shall be further classified into following categories:

7.3.1. CRZ-III A

Such densely populated CRZ-III areas, where the population density is more than 2161 per square kilometer as per 2011 census base, shall be designated as CRZ-III A and in CRZ-III A, area up to 50 meters from the HTL on the landward side shall be earmarked as the 'No Development Zone (NDZ)', provided the CZMP as per this notification, framed with due consultative process, have been approved, failing which, a NDZ of 200 meters shall continue to apply.

7.3.2. CRZ-III B

All other CRZ-III areas with population density of less than 2161 per square kilometer, as per 2011 census base, shall be designated as CRZ-III B and in CRZ-III B, the area up to 200 meters from the HTL on the landward side shall be earmarked as the 'No Development Zone (NDZ)'.

7.4. CRZ-IV

The CRZ- IV constitutes the water area and shall be further classified as under: -

7.4.1. CRZ- IVA

The water area and the seabed area between the Low Tide Line up to twelve nautical miles on the seaward side shall constitute CRZ-IV A.

7.4.2. CRZ- IVB

CRZ-IV B areas shall include the water area and the bed area between LTL at the bank of the tidal influenced water body to the LTL on the opposite side of the bank, extending from the mouth of the water body at the sea up to the influence of tide, i.e., salinity of five parts per thousand (ppt) during the driest season of the year.

7.5. Regulation limits/lines

The CRZ limits has been revised or updated as per the provisions contained in the CRZ Notification 2019. The 50 meters No Development Zone (NDZ) in the case of CRZ-III areas/ a 50 meters buffer line (CRZ limit) in the case of CRZ-II areas or the width of the creeks that are influenced by tidal from sea in the bays, estuaries, rivers, creeks, backwaters, lagoons, ponds. have been drawn. The landward extent of NDZ/Buffer by the creek is up to the location where the salinity limit of 5 part per thousand is encountered.

By the seacoast, a 500-meter line from HTL is drawn as CRZ limit irrespective of whether the area is under CRZ-III or CRZ-II. As per the CRZ Notification 2019, a new sub-category of CRZ- IIIA by the coast is introduced, where the NDZ is limited to 50 meters and the same has been updated. Similarly, as in the previous CZMP, the 200-meter line of NDZ by the seacoast has been drawn for the CRZ-IIIB areas.

The NDZ of the islands in the coastal backwaters as well as islands along the mainland coast has been limited to 20 meters, the same has been updated in the CZMP being prepared according to the CRZ Notification 2019, but it will be implemented only after the approval of IIMP of the

particular Island.

Subject to the information to be provided by the State Government on the details of village-wise survey numbers pertaining to government land for deciding/enabling ease in demarcation of buffers around mangrove areas, a uniform buffer of 50 meters have been demarcated in case of mangrove area being more than 1000 square meters. The ownership details of Mangroves more than 1000sq.m spread area provided by Govt. of Kerala is provided in the **Annexure 4**.

7.6. CVCA and IIMP

Critically Vulnerable Coastal Areas (CVCA)

Sundarban region of West Bengal and other ecologically sensitive areas identified as under Environment (Protection) Act, 1986 such as Gulf of Khambat and Gulf of Kutchh in Gujarat, Malvan, Achra-Ratnagiri in Maharashtra, Karwar and Coondapur in Karnataka, Vembanad in Kerala, Gulf of Mannar in Tamil Nadu, Bhitarkanika in Odisha, Coringa, East Godavari and Krishna in Andhra Pradesh shall be treated as Critical Vulnerable Coastal Areas (CVCA) and managed with the involvement of coastal communities including fisher folk who depend on coastal resources for their sustainable livelihood.

No CVCA is marked in the Thrissur district and in Kerala, except certain parts around Vembanad Lake, no other areas have been marked as CVCA by the NCSCM.

Integrated Island Management Plan (IIMP)

The islands demarcated in Thrissur District as mainland coast islands and inland islands in the coastal backwaters need to have Integrated Island Management Plans (IIMPs), as applicable to smaller islands in Lakshadweep and Andaman & Nicobar, as per Island Protection Zone Notification, 2011 number S.O. 20(E), dated the 6th January, 2011 to be formulated by the Kerala State. This would be carried out with the help of NCSCM, once all such islands marked in this CZMP are approved by the Ministry of Environment, Forest and Climate Change. There are fifty-three number of backwater islands are present in Thrissur district. The islands are buffered with 50 m or width of the creek whichever is less and 20m CRZ line landward of HTL is also depicted in the map which

will be considered only after IIMP is implemented. The dimensions of backwater islands show wide variation in the district, ranging area from 0.000374 Km² (Pavaratty Panchayat) to 5.8803 Km² (Kodungallur Municipality). All together the area of islands accounts for 12.35068 Km² (Detailed table is annexed in Annexure 2).

8. HAZARD LINE

A 'Hazard line' has been demarcated by the Survey of India (SOI) taking into account the extent of the flooding on the land area due to water level fluctuations, sea level rise and shoreline changes (erosion or accretion) occurring over a period of time. The hazard line mapped by SOI has been shared by NCSCM as part of the previous CZMP prepared. The hazard line is to be used as a tool for disaster management plan for the coastal environment, including planning of adaptive and mitigation measures. With a view to reduce the vulnerability of the coastal communities and ensuring sustainable livelihood, while drawing the CZMP, the land use planning for the area between the Hazard line and HTL need to be take into account as such impacts of climate change and shoreline changes.

9. CRZ CATEGORIES OF THRISSUR DISTRICT

The CRZ of the Thrissur district consist of CRZ-IA, CRZ-IB/CRZ-IB Pokkali, CRZ-II, CRZ-III A, CRZ-III B, CRZ-IV A and CRZ-IV B. Altogether 31 local bodies are covered under the CRZ area of which 28 are Grama Panchayaths and 3 are Municipal Council areas. Altogether 51 villages are under the purview of CRZ in Thrissur District. The details are provided in the table annexed (Annexure-2).

The new village and panchayat boundaries provided by KCZMA, obtained from Survey and Land Records does not match with the survey plots and district boundaries in the approved CZMP, 2011. Hence old boundaries (approved CZMP, 2011) are used for this exercise, as per the instruction from DoECC, Govt. of Kerala. However, as per the new village boundary Chendrapinni village has been renamed as Edathiruthi Chenthrapinny village and Vadama village has been renamed as Vadama Vadakkumbhagom village.

CRZ categories and ESAs (Panchayat/Village-wise) in Thrissur District

A detailed table is annexed (Annexure-2) along with this report separately on the Panchayath/Village-wise statistics of HTL, ESAs, intertidal area, mangrove buffer, area covered under each CRZ category.

10. CONCLUSION

Statistics of the CRZ status of Thrissur District is summarized below:	
Total length of HTL along the Seacoast	58.32 Km
Total length of HTL along the inland water bodies	509.93 Km
Total Area under the Archeological/Historical sites	1.90 Km ²
Total Area under the Turtle Nesting Sites	0.043373 Km ²
Total area under mangrove extent	1.315048 Km ²
Total area under mangrove buffer	0.37722 Km ²
Total area under intertidal zone (CRZ-IB)	11.77028 Km ²
Total area under intertidal zone (CRZ-IB) Pokkali	0.644082 Km ²
Total area under CRZ-II along the Seacoast	2.422771 Km ²
Total area under CRZ-II along the inland water bodies	3.345077 Km ²
Total area in No Development Zone in CRZ-III along Seacoast	8.418341 Km ²
Total area in No Development Zone in CRZ-III along water bodes	14.21063 Km ²
Total area in CRZ-IIIA along the coast between 50-500 meters	11.13841 Km ²
Total area in CRZ-IIIB along the coast between 200-500 meters	7.504515 Km ²
Total area under the CRZ-IVB category	21.17689 Km ²